

IN THE CLAIMS

PLEASE AMEND THE CLAIMS AS FOLLOWS:

1. (CANCELLED)

2. (CANCELLED)

3. (CANCELLED)

4. (CANCELLED)

5. (CURRENTLY AMENDED) ~~The method of claim 1~~ A method for indicating viability of transplanted progenitor or stem cells grown in a culture, the method being performed with a medical device that supports at least one sensing function, the method comprising:
non-destructively observing a region of a patient to where progenitor or stem cells grown in a culture cells have been transplanted;
sensing a property within said region of a patient that is indicative of cell viability or inviability of transplanted progenitor or stem cells grown in a culture; and
using data from sensing said property within said region to indicate cell viability from a transplant of progenitor or stem cells grown in a culture within the region wherein said cell viability is indicated by a property in cell chemistry resulting from an event selected from the group consisting of cell activity, cell inactivity, cell growth, cell death, specific cell function, specific cell dysfunction, volumetric expansion of cell population, and volumetric decrease of cell population.

6. (CURRENTLY AMENDED) ~~The method of claim 2~~ A method for indicating viability of transplanted progenitor or stem cells grown in a culture, the method being performed with a medical device that supports at least one sensing function, the method comprising:
non-destructively observing a region of a patient to where progenitor or stem cells grown in a culture cells have been transplanted;
sensing a property within said region of a patient that is indicative of cell viability or inviability of transplanted progenitor or stem cells grown in a culture; and
using data from sensing said property within said region to indicate cell viability from a

transplant of progenitor or stem cells grown in a culture within the region, wherein said non-destructively observing comprises magnetic resonance imaging and wherein said cell viability is indicated by a property in cell chemistry resulting from an event selected from the group consisting of cell activity, cell inactivity, cell growth, cell death, specific cell function, specific cell dysfunction, volumetric expansion of cell population, and volumetric decrease of cell population.

7. (CURRENTLY AMENDED) ~~The method of claim 1~~ A method for indicating viability of transplanted progenitor or stem cells grown in a culture, the method being performed with a medical device that supports at least one sensing function, the method comprising:

non-destructively observing a region of a patient to where progenitor or stem cells grown in a culture cells have been transplanted;

sensing a property within said region of a patient that is indicative of cell viability or inviability of transplanted progenitor or stem cells grown in a culture; and

using data from sensing said property within said region to indicate cell viability from a transplant of progenitor or stem cells grown in a culture within the region wherein said property is monitored by observation of at least one parameter selected from the group consisting of local lactate levels, local glucose turnover, local phosphorous high-energy metabolite concentrations, local F-19 labeled metabolites, alterations in tissue sodium, and changes in the conversion rates of O₂ gas to H₂O water.

8. CANCELLED

9. (PREVIOUSLY PRESENTED) The method of claim 6 wherein said property is monitored by observation of at least one parameter selected from the group consisting of local lactate levels, local glucose turnover, local phosphorous high-energy metabolite concentrations, local F-19 labeled metabolites, alterations in tissue sodium, and changes in the conversion rates of O₂ gas to H₂O water.

10. CANCELLED

11. (CURRENTLY AMENDED) The method of claim 5 2 wherein said property is monitored by at least one technique selected from the group consisting of proton spectroscopy, monitoring of C-13 labeled glucose, monitoring by P-31 MR spectroscopy, monitoring of local F-19 labeled metabolites, monitoring of Na-23 levels, and monitoring of $^{17}\text{O}_2$ gas conversion to H_2^{17}O water.

12. (ORIGINAL) The method of claim 6 wherein said property is monitored by at least one technique selected from the group consisting of proton spectroscopy, monitoring of C-13 labeled glucose, monitoring by P-31 MR spectroscopy, monitoring of local F-19 labeled metabolites, monitoring of Na-23 levels, and monitoring of $^{17}\text{O}_2$ gas conversion to H_2^{17}O water.

13. (CURRENTLY AMENDED) The method of claim 5 4 wherein said medical device includes at least one element selected from the group consisting of a volume coil surrounding the tissue and a local multi-tuned MRI RF coil.

14. (CURRENTLY AMENDED) The method of claim 6 2 wherein said medical device includes at least one element selected from the group consisting of a volume coil surrounding the tissue and a local multi-tuned MRI RF coil.

15. (ORIGINAL) The method of claim 9 wherein said medical device includes at least one element selected from the group consisting of a volume coil surrounding the tissue and a local multi-tuned MRI RF coil.

16. (ORIGINAL) The method of claim 12 wherein said medical device includes at least one element selected from the group consisting of a volume coil surrounding the tissue and a local multi-tuned MRI RF coil.

17. (PRESENTLY AMENDED) ~~The method of claim 1~~ A method for indicating viability of transplanted progenitor or stem cells grown in a culture, the method being performed with a medical device that supports at least one sensing function, the method

comprising:

non-destructively observing a region of a patient to where progenitor or stem cells grown in a culture cells have been transplanted;

sensing a property within said region of a patient that is indicative of cell viability or inviability of transplanted progenitor or stem cells grown in a culture; and
using data from sensing said property within said region to indicate cell viability from a transplant of progenitor or stem cells grown in a culture within the region wherein said property comprises blood flow or changes in blood flow as vascular supply is developed.

18. (CURRENTLY AMENDED) The method of claim 17 2 wherein said non-destructively observing comprises magnetic resonance imaging and said property comprises blood flow or changes in blood flow as vascular supply is developed.

19. (CURRENTLY AMENDED) The method of claim 17 7 wherein said property is monitored by observation of at least one parameter selected from the group consisting of local lactate levels, local glucose turnover, local phosphorous high-energy metabolite concentrations, local F-19 labeled metabolites, alterations in tissue sodium, and changes in the conversion rates of O₂ gas to H₂O water said property comprises blood flow or changes in blood flow as vascular supply is developed.

20. (ORIGINAL) The method of claim 17 wherein blood flow or changes in blood flow are measured by observation of at least one material selected from the group consisting of labeled H₂O water, contrast-agent infusion of T1-shortening agents or T2*-shortening agents, local introduction of hyperpolarized Xenon gas, or optically-active coloring agents.

21. (ORIGINAL) The method of claim 18 wherein blood flow or changes in blood flow are measured by observation of at least one material selected from the group consisting of labeled H₂O water, contrast-agent infusion of T1-shortening agents or T2*-shortening agents, local introduction of hyperpolarized Xenon gas, or optically-active coloring agents.

22. (ORIGINAL) The method of claim 19 wherein blood flow or changes in blood flow are measured by observation of at least one material selected from the group consisting of labeled H₂O water, contrast-agent infusion of T1-shortening agents or T2*-shortening agents, local introduction of hyperpolarized Xenon gas, or optically-active coloring agents.

23. (CURRENTLY AMENDED) ~~The method of claim 2~~ A method for indicating viability of transplanted progenitor or stem cells grown in a culture, the method being performed with a medical device that supports at least one sensing function, the method comprising:

non-destructively observing a region of a patient to where progenitor or stem cells grown in a culture cells have been transplanted;

sensing a property within said region of a patient that is indicative of cell viability or inviability of transplanted progenitor or stem cells grown in a culture; and using data from sensing said property within said region to indicate cell viability from a transplant of progenitor or stem cells grown in a culture within the region wherein said non-destructively observing comprises magnetic resonance imaging and said property comprises anisotropic water diffusion.

24. (CURRENTLY AMENDED) The method of claim 5 ~~2~~ wherein said property comprises the local concentrations of at least one of choline, NAA, GABA, phosphocholine, and creatine.

25. (CURRENTLY AMENDED) The method of claim 6 ~~4~~ wherein the property is selected from the group consisting of a) local tissue density and cell populations, b) local electrical activity, c) local oxygenated/deoxygenated hemoglobin and changes in the local T2* reflecting the alterations in tissue oxygenation, d) changes in the vascular reserve and response to oxygenation stresses, e) tissue fluorescence and bioluminescence, f) tissue fluorescence and bioluminescence, g) electrical impedance, and h) local tissue temperature.

26. (ORIGINAL) The method of claim 1 wherein the property is selected from the group consisting of a) local tissue density and cell populations, b) local electrical activity, c) local oxygenated/deoxygenated hemoglobin and changes in the local T2* reflecting the alterations in tissue oxygenation, d) changes in the vascular reserve and response to oxygenation stresses, e) tissue fluorescence and bioluminescence, f) tissue fluorescence and bioluminescence, g) electrical impedance, and h) local tissue temperature.

27. (CANCELLED)

28. (CANCELLED)

29. (CURRENTLY AMENDED)~~The method of claim 28~~ A method for indicating viability of transplanted progenitor or stem cells grown in a culture, said method being performed with a medical device that supports at least one sensing function comprising:
non-destructively observing a region of a patient to where progenitor or stem cells grown in a culture have been transplanted;
sensing a property within said region of a patient that is indicative of cell metabolism;
repeating or continuing said sensing of a property over a period of time in which said property changes; and
using data from sensing changes in said property within said region to indicate cell viability from a transplant of progenitor or stem cells grown in a culture within the region, wherein said data from sensing changes in said property indicates active metabolic function in transplanted cells, and wherein changes in said property are monitored by at least one technique selected from the group consisting of proton spectroscopy, monitoring of C-13 labeled glucose, monitoring by P-31 MR spectroscopy, monitoring of local F-19 labeled metabolites, monitoring of Na-23 levels, and monitoring of $^{17}\text{O}_2$ gas conversion to H_2^{17}O water.

30. (CANCELLED)

31. (CANCELLED)

32. (CANCELLED)
33. (CANCELLED)
34. (CANCELLED)
35. (CANCELLED)
36. (CANCELLED)
37. (CANCELLED)
38. (CANCELLED)
39. (CANCELLED)
40. (CANCELLED)
41. (CANCELLED)
42. (CANCELLED)
43. (CANCELLED)
44. (CANCELLED)
45. (CANCELLED)
46. (CANCELLED)
47. (CANCELLED)
48. (CANCELLED)
49. (CANCELLED)
50. (CANCELLED)
51. (CANCELLED)
52. (CANCELLED)
53. (CANCELLED)